US ISSN 0046 225X

ENVIRONMENTAL ENTOMOLOGY



Volume 19, 1990
1–1924

Environmental Entomology

Volume 19

WILLIAM P. KEMP
Editor
RONALD E. STINNER
Editor
JOHN D. BRIGGS
Book Review Editor

RAYMOND L. EVERNGAM, JR.
Managing Editor
RICHARD H. FOOTE
ELIZABETH A. MITCHELL
CHRISTINA A. DAVIS
Copy Editors

JACQUELINE L. HAIGHT
DEBORAH L. SANDERS
STACEY N. WAGES
Editorial Assistants

Editorial Board

D. W. JOHNSON, *Chairman*
Princeton, Ky.
Section E (1990)

L. A. HULL
Biglerville, Pa.
Section F (1991)

T. O. HOLTZER
Fort Collins, Colo.
Section C (1992)

H. R. BURKE
College Station, Tex.
Section A (1992)

C. J. JONES
Urbana, Ill.
Section D (1993)

C. LOUDON
Ithaca N.Y.
Section B (1994)

Published by the
Entomological Society of America

J. H. OLIVER, JR.
Statesboro, Ga.
President

D. FEIR
St. Louis, Mo.
Past President

L. R. NAULT
Wooster, Ohio
President-Elect

J. H. SHADDY
Kirksville, Mo.
Secretary–Treasurer

E. A. CAMERON
University Park, Pa.
Eastern Branch

E. E. ORTMAN
West Lafayette, Ind.
North Central Branch

R. E. BERRY
Corvallis, Oreg.
Pacific Branch

B. R. WISEMAN
Tifton, Ga.
Southeastern Branch

D. R. RUMMEL
Lubbock, Tex.
Southwestern Branch

B. J. HARRINGTON
Madison, Wis.
Section A

G. M. CHIPPENDALE
Columbia, Mo.
Section B

R. T. CARDÉ
Amherst, Mass.
Section C

J. D. EDMAN
Amherst, Mass.
Section D

D. D. KOPP
Fargo, N.D.
Section E

G. G. KENNEDY
Raleigh, N.C.
Section F

W. D. HANSEN
Lanham, Md.
Executive Director

© 1990, Entomological Society of America

Environmental Entomology (ISSN 0046-225X) is published bimonthly (February, April, June, August, October, and December) by the Entomological Society of America, 9301 Annapolis Road, Lanham, Md. 20706. The journal is printed by Allen Press, Inc., 1041 New Hampshire Street, P.O. Box 368, Lawrence, Kans. 66044. Second class postage is paid at Lanham, Md., and at additional mailing offices.

Postmaster: Send change of address forms to *Environmental Entomology*, ESA, 9301 Annapolis Road, Lanham, Md. 20706.

Inquiries regarding content, subscriptions, advertising, availability of back issues, change of address, and general business matters, and manuscripts and other copy should be sent to the managing editor, ESA, 9301 Annapolis Road, Lanham, Md. 20706. For information about the preparation of manuscripts, authors are urged to consult "Information for Contributors," which is printed on the inside back cover of most issues, or to send for *Publications Policies and Guide for Authors*, which is available free of charge from ESA.

Articles report the results of research only. Mention of a pesticide or a commercial or proprietary product does not constitute endorsement or recommendation for its use. Authors are entirely responsible for statements made, whether of fact or opinion. Registered names and trademarks, etc., used in this publication, even without specific indication thereof, are not to be considered unprotected by law.

Photocopies. Permission to photocopy journal articles for internal or personal use is granted by the Entomological Society of America provided that a fee is paid directly to the Copyright Clearance Center, 21 Congress Street, Salem, Mass. 01970. Individuals may make a single copy for personal use without charge. This consent does not extend to other kinds of copying, such as that for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale. The fee code for users of the Transaction Reporting Service is 0046-225X/89/$2.00.

Subscriptions. 1989 prices: members, $25; nonmember individuals, $65; institutions, $110. To these prices add $18 for shipping outside the continental U.S.; $20 for shipping to India. Send payment to EE Subscriptions, ESA, P.O. Box 177, Hyattsville, Md. 20781. Inquiries should be directed to ESA Sales, 9301 Annapolis Road, Lanham, Md. 20706; (301) 731-4535.

Back Issues	ESA member domestic	ESA member foreign	Non-member domestic	Non-member foreign
Single issue	$10.40	$11.40	$ 34.00	$ 35.00
Volume	62.40	68.40	204.00	210.00

All prices include handling and shipping by surface mail. For a list of available issues, write to ESA Sales, 9301 Annapolis Road, Lanham, Md. 20706. Back volumes are available in microform from University Microfilms International, 300 North Zeeb Road, Ann Arbor, Mich. 48106.

CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 19, 1990

A
Abrahamson, L. P., 257
Abrahamson, W. G., 465
Ali, A., 1060
Allen, W. R., 309
Appel, A. G., 15
Arnold, G., 1780
Awan, M. S., 387
Axtell, R. C., 578

B
Bach, C. E., 987
Baines, D., 1146
Baker, T. C., 21, 490
Balaam, R. J., 943
Balciunas, J. K., 799
Baranowski, R. M., 378
Barbosa, P., 640
Barrett, B. A., 362, 803
Barrett, G. W., 955
Barry, B. D., 1091
Bartelt, R. J., 182
Basimike, M., 486
Bastian, R. A., 409
Bates, M. B., 1675
Bauduin, F., 222
Bauer, L. S., 428
Bellinger, R. G., 1024
Berberet, R. C., 1001
Berisford, C. W., 99, 1671
Berisford, Y. C., 1671
Bessin, R. T., 635
Bing, J. W., 932
Bishop, L., 1738
Black, D. A., 182
Blackman, R. L., 609
Blood, P. R. B., 710
Boethel, D. J., 746
Bohlen, P. J., 955
Bohls, R. A., 1746
Boivin, G., 836, 1146
Boldt, P. E., 515
Bollich, C. N., 268
Bosio, C. F., 465
Bossart, J. L., 1055
Boucias, D. G., 137
Brown, S., 195
Brown, S. K., 565
Browning, H. W., 345, 393
Brunner, J. F., 362, 803
Brust, G. E., 697, 966, 1837
Brusven, M. A., 1786
Bullock, R. H., 943
Buntin, G. D., 603
Byrne, P. F., 1091

C
Caprio, M. A., 281
Carey, J. R., 1075
Carlson, R. E., 1773
Carroll, S., 91
Carroll, T. W., 609
Carruthers, R. I., 85
Center, T. D., 799
Chan, K.-L., 286
Chianese, R. J., 943
Chon, T.-S., 1716
Clarke, S. R., 99
Clausen, K., 1689
Clearwater, J. R., 1702
Clement, S. L., 1882
Cloutier, C., 222, 293
Coderre, D., 1657
Cofrancesco, A. F., 799
Cogburn, R. R., 268
Cohen, A. C., 327
Cokendolpher, J. C., 878
Collins, R. D., 949
Cotterill, B., 195
Crawford, A. M., 1115
Cuda, J. P., 415
Culin, J., 195
Cummins, J. N., 565
Czapowskyj, M. M., 257

D
Daane, K. M., 1130
Dahlsten, D. L., 1130
Daigle, C. J., 746
Daniels, T. J., 1029
Darrah, L. L., 1091
Day, J. F., 469
DeBarr, G. L., 99
DeLoach, C. J., 415
Delgado-Salinas, A., 1017
Dennehy, T. J., 764
Dennill, G. B., 130
Derrick, M. E., 1081
Desenfant, PH., 1780
Dickson, M. H., 1086
Dimmick, C. R., 972
Dodge, K. L., 905
Domek, J. M., 995
Dong, K., 1722
Dray, Jr., F. A., 799
Dreistadt, S. H., 1130
Dunn, J. P., 239

E
Eigenbrode, S. D., 1086
Elkinton, J. S., 662, 859
Ellis, C. R., 666
Endress, A. G., 854
Eskafi, F. M., 148, 1666

F
Faulder, R. J., 1710
Felland, C. M., 162
Feng, M.-G., 785, 815
Fernandez, A., 1666
Ferro, D. N., 104, 117, 123, 1732, 1858
Fielding, D. J., 1786
Fischer, D., 1852
Fish, D., 1029
Flanagan, G. J., 776
Flinn, P. W., 176, 725
Flint, H. M., 1845
Foster, J. E., 494
Foster, R. N., 1746
Freeman, R., 1075
Freytag, P. H., 209
Fried, C. S., 322
Frisbie, R. E., 91
Fuester, R. W., 943
Funderburk, J. E., 137
Funk, C. R., 1037
Fuxa, J. R., 260, 746

G
Gabriel, A. D., 339, 932
Gage, S. H., 370, 524, 534, 1055
Gahukar, R. T., 558
Garcia, J. A., 403
Garris, G. I., 66
Geden, C. J., 578
Geden, C. J., 1722
Gillett, J. D., 776
Gillott, C., 587, 704
Gilreath, M. E., 137
Gilstrap, F. E., 684
Gordon, A. J., 130
Gould, J. R., 859
Grafius, E., 890
Grafius, E. J., 281
Gregory, Jr., B. M., 1017
Grimes, D. W., 1803
Gross, H. R., 1122
Grover, Jr., P. B., 494
Guppy, J. C., 423, 612
Gutierrez, A. P., 1767

H
Habeck, D. H., 799
Hagan, D. V., 1068
Hagler, J. R., 327, 1034
Hagley, E. A. C., 309
Hagstrum, D. W., 725
Hajek, A. E., 85
Halbert, S. E., 609
Hammond, Jr., A. M., 1017
Hammond, R. B., 510, 1810
Harcourt, D. G., 612
Harman, A. L., 501
Harman, D. M., 501
Harris, M. O., 303
Hart, E. R., 409
Hart, W. G., 403
Hassan, S. T. S., 710
Hayakawa, D. L., 890
Hebert, C., 293
Heinrichs, E. A., 167, 594
Heinz, K. M., 825
Heliovaara, K., 481
Hemming, J. D. C., 842
Herbert, D. A., 1135
Herren, H. R., 1767
Higbee, B. S., 544
Hilton, R. J., 753
Hoffmann, J. H., 1866
Hoffmann, M. P., 387, 753

Horton, D. R., 357, 573, 656
Hostetter, D. L., 1150
Hough-Goldstein, J. A., 234
House, G. J., 697, 966
Howell, J. F., 573
Hower, A. A., 176
Hoxie, R. P., 1698
Hoy, J. B., 898
Hull, L. A., 648
Hulme, M. A., 54
Hunsberger, A. G. B., 313
Hunt, R. E., 76

I
Idoine, K., 104, 1732

J
James, D. G., 874, 1710
Jang, E. B., 1726
Jansson, R. K., 313
Jarvis, B. J., 587, 704
Jean, C., 1657
Jeffers, D. L., 1810
Jennings, D. T., 257
Johnson, D. L., 1103
Johnson, D. T., 995
Johnson, J. B., 785, 815
Johnson, M. P., 677
Johnson, S. J., 740, 978, 1017
Johnson-Cicalese, J. M., 1037
Jones, C. G., 108
Jones, S. C., 1047

K
Kays, S. J., 1663
Keaster, A. J., 1091
Kemp, W. P., 459
Kepner, R. L., 1746
Kettunen, J., 905
Khattak, S. U. K., 573
Kimmerer, T. W., 239
Kish, L. P., 815
Kline, D. L., 1068
Klubertanz, T. H., 1773
Knievel, D. P., 176
Knight, A. L., 648
Knight, A. W., 1681
Knutson, A. E., 684
Kogan, M., 854, 1852
Kolbe, M. M., 148
Kosalwat, P., 1681
Kovach, J., 195
Kramer, D. A., 578
Kring, T. J., 154, 1798
Krysan, J. L., 544, 551
Kuenen, L. P. S., 1

L
Lamb, R. C., 565
Lampman, R. L., 8
Lance, D. R., 882
Le Conte, Y., 1780
Lecrone, S. H., 313
Lee, J.-H., 740, 978
Letourneau, D. K., 505
Lewis, P. A., 8
Liang, D., 1716
Lichter, J. P, 972
Lievens, A. W., 1017
Lin, H., 854, 1852
Lindegren, J. E., 383
Lindroth, R. L., 842
Linley, J. R., 286, 1068
Livne, H., 1010
Loera, J., 274
Lofgren, C. S., 44
Lolong, Z., 1115
Loper, G. M., 327
Luciano, P., 108
Luttrell, R. G., 1060
Lynch, R. E., 137

M
MacDonald, P. J., 666
MacKay, P. A., 618
MacMahon, J. A., 244
Mack, T. P., 229
Mackay, I. F., 911
Magnon, G. J., 1068
Mahr, D. L., 1097
Maier, K. J., 1681
Majchrowicz, I., 1163
Makela, M. E., 91
Mamood, A. N., 1034
Mangan, F. X., 1858
Mangan, R. L., 808
Maniania, N. K., 1163
Margolies, D. C., 949
Marsh, Jr., H. V., 1858
Marti, O. G., 795
Mayse, M. A., 1689
McAuslane, H. J., 26
McCauley, G. W., 949
McClain, D. C., 916, 926
McClure, M. S., 36, 672
McConnell, J. A., 1798
McCrea, K. D., 465
McDowell, J. M., 137
McGuire, M. R., 182
McInnis, D. O., 383
McIntosh, A. H., 1150
McManus, M. L., 1024
McPherson, R. M., 1135
Meisch, M. V., 961
Meloche, F., 423
Mendel, Z., 1010
Metcalf, R. L., 8
Metterhouse, W. W., 943
Minkenberg, O. P. J. M., 625
Mitchell, F. L., 260
Mittler, T. E., 32
Montgomery, M. E., 937
Moore, K. E. B., 108
Moran, V. C., 1866
Moratorio, M. S., 142
Morrison, R. K., 44
Moss, T., 1135
Mueller, A. J., 677
Mullin, C. A., 474
Murray, K. D., 662
Mutinga, M. J., 486

N
Nault, L. R., 76
Neupane, F. P., 215
Nielsen, D. G., 937
Nishida, T., 1820
Nishijima, K. A., 1726
Nitao, J. K., 465
Norris, D. M., 215
Nyambo, B. T., 1141
Nyomura, K., 1845
Nyrop, J. P., 764

O
O'Hair, S. K., 313
O'Keeffe, L. E., 630
O'Neill, K. M., 459
ODell, T. M., 771
Obrycki, J. J., 339, 932
Okech, S.H.O., 1792
Orr, C. J., 932
Oseto, C. Y., 1675
Overholt, W. A., 1155, 1889

P
Pair, S. D., 274
Parks, N. J., 1845
Parrella, M. P., 835
Payne, T. L., 1671
Peacock, J. W., 1
Pedigo, L. P., 603, 1773
Pena, J. E., 378
Phillips, Jr., S. A., 878
Pimentel, D., 322
Pinnell, R. E., 1150
Pitre, H. N., 1060
Pivnick, K. A., 587, 704
Podoler, H., 1010
Pooler, P. D., 1746
Popham, T. W., 66
Poprawski, T. J., 1163
Poswal, M. A., 1001
Potter, D. A., 239
Price, J. R., 1110
Price, P. W., 905
Purcell, M., 848
Puterka, G. J., 1110
Puttler, B., 1150

Q
Quinn, M. A., 1746, 1756

R
Raffa, K. F., 1829
Rajotte, E. G., 648
Ramsey, A. M., 469
Rapusas, H. R., 167, 594
Raulston, J. R., 274
Raupp, M. J., 771
Ravlin, F. W., 1024
Reagan, T. E., 635

Reardon, R. C., 943
Reese, J. C., 949
Regniere, J., 293
Reilly, C. C., 44
Reissig, W. H., 565
Reuter, K. C., 1746
Ridgway, N. M., 1097
Riechert, S. E., 1738
Rieske, L. K, 1829
Robacker, D. C., 403
Robbins, T. O., 415, 515
Robert, P.-H., 1163
Robison, D. J., 257
Rock, G. C., 916, 926
Rodriguez-Del-Bosque, L. A., 345, 393
Rogers, C. E., 795
Rogers, J., 195
Roland, J., 721
Roltsch, W. J., 524, 534, 1689
Rose, S., 303
Roseland, C. R., 1675
Rummel, D. R., 91
Rust, R. W., 332
Rutz, D. A., 322, 1722
Ryan, R. B., 1873

S
Saxena, K. N., 1792
Scarborough, D., 195
Schaalje, G. B., 439
Schal, C., 1716
Schmidt, R. S., 573
Schotzko, D. J., 630
Scott, J. G., 1722
Seaman, A. J., 764
Sedlacek, J. D., 209
Segarra-Carmona, A., 640
Severson, R. F., 1663
Shelton, A. M., 111, 1086
Shields, E. J., 866
Shorthouse, J. D., 911
Showers, W. B., 603, 1081
Shukle, R. H., 494
Siegfried, B. D., 474
Silk, P. J., 1
Silvain, J. F., 795
Simmons, A. M., 795
Simmons, G. A., 370
Simpson, K. B., 1091
Slater, G. P., 587, 704
Slosser, J. E., 1110
Smith, Jr., J. W., 345, 393, 1155, 1889
Smith, M. A. H., 618
Son, K.-C., 1663
Soper, R. S., 85
Spangler, S. M., 244
Sparks, A. N., 274
Speese III, J., 1135
Sponsler, R. C., 15
Spurgeon, D. W., 229
Stark, J. D., 1820
Stehr, F. W., 890
Steininger, M. K., 108
Sterling, W. L., 59
Stevenson, A. B., 836
Steward, V. B., 154
Stewart, R., 1146
Stinner, R. E., 578, 916
Stone, N. D., 91
Stoner, K. A., 730
Sturm, M. M., 59
Suckling, D. M., 1702
Summy, K. R., 274
Swift, A., 195

T
Tabashnik, B. E., 987
Tahvanainen, J., 905
Talhouk, S. N., 937
Taylor, P. L., 1698
Taylor, P. S., 866
Tebbets, J. S., 791
Tedders, W. L., 44
Thomas, D. B., 808
Thompson, C. R., 799
Thorpe, K. W., 771
Tisdale, R. A., 456
Torchio, P. F., 1649
Tourneur, J.-C., 1657
Trichilo, P. J., 1803
Tyrrell, D., 612

U
Underhill, E. W., 587, 704
Underhill, L. G., 1866

V
Vail, P. V., 791
Vaisanen, R., 481
Van Den Berg, H., 1141
Vargas, R. I., 1820
Vetter, R. S., 21
Vinson, S. B., 26
Voss, R. H., 117, 123

W
Waage, J. K., 1141
Wagner, M. R., 456
Walgenbach, D. D., 1746, 1756
Waller, G. D., 1034
Warren, G. N., 1710
Weathersbee III, A. A., 961
Webb, S. E., 111
Weber, D. C., 1858
Weis, A. E., 972
Welch, J. B., 808
Wellso, S. G., 1698
Welter, S. C., 848
Weseloh, R. M., 448
White, E. H., 257
White, L. D., 573
Wiesenborn, W. D., 490
Wilhoit, L., 32
Williams, D. W., 943
Williams, H. J., 26
Willms, W. D., 1103
Wilson, C. G., 776
Wilson, D. D., 1663
Wilson, L. T., 387, 710, 753, 1803
Wirth, T. M., 370
Wolfe, G. W., 1037
Wong, T. T., 383
Wood, B. W., 44
Woolley, J. B., 926
Wright, S. S., 1
Wright, V. L., 978

Y
Yamamoto, A., 1845
Yaninek, J. S., 1767
Yeargan, K. V., 209
Yearian, W. C., 154, 251
Young, L. J., 1001
Young, S. Y., 251, 717

Z
Zalom, F. G., 753
Zehnder, G. W., 1135
Zelazny, B., 1115
Zhang, J.-T., 469
Zimmerman, R. H., 66
Zmijowskyj, T. J., 911
Zuber, M. S., 1091

CUMULATIVE SUBJECT INDEX
Environmental Entomology
Volume 19, 1990

A
Acalymma vittatum, attractants, 8
  insect–plant interactions, 8
Acigona ignefusalis, cultural control, 558
  population ecology, 558
Acrididae, communities, 1756
  density correlation, 1786
  grazing history, 1756
  weather, 1786
action site, flush bar technique, 1081
  Ostrinia nubilalis, 1081
Acyrthosiphon kondoi, life table, 1001
  natural control, 1001
Acyrthosiphon pisum, latitudinal variation, 618
  photoperiod, 618
adaptation, corn, 949
  Schizaphis graminum, 949
Adelges tsugae, dispersal agents, 36
  insect dispersal, 36
agonistic behavior, subterranean termites, 1047
  territory size, 1047
Agrilus bilineatus, *Cryphonectria (Endothia) parasitica*, 239
  tree stress, 239
Agromyza frontella, *Dacnusa dryas*, 423
  encapsulation, 423
alfalfa, soybean, 677
  Spissistilus festinus, 677
Allium cepa, *Apis mellifera*, 327
  onion nectar, 327
Anastrepha ludens, attractants, 403
  Sargentia greggii, 403
Ancylis comptana, diapause, 932
  dormancy, 932
  integrated pest management, 339
  temperature–development relationship, 339
Anopheles quadrimaculatus, dispersal, 961
  flight range, 961
Anthomyiidae, conservation tillage, 510
  Glycine max, 510
Anthonomus grandis grandis, cotton, 1110
  diapause, 91
  dispersal, 195
  modeling, 195
  mortality, 59, 91
  parasitism, 59
  parasitoids, 1110
Anticarsia gemmatalis, degree-day survival, 1017
  microhabitat distribution, 740
  models, 1017
  parasitoids, 746
  Pseudoplusia includens, 717, 746
  Solenopsis invicta, 978
  soybeans, 740, 978
  sprinkler irrigation, 717
antifeedants, herbs, 234
  Leptinotarsa decemlineata, 234
aphid–pathogenic fungi, *Beauveria bassiana*, 815
  bioassay, 815
Aphidius liriodendrii, biological control, 1130
  Trioxys curvicaudus, 1130
Aphidius nigripes, biological control, 222
  foraging, 222
Apis mellifera, *Allium cepa*, 327
  Gossypium spp., 1034
  onion nectar, 327
  pollen deposition, 1034
  temperatures, 1780
  Varroa jacobsoni, 1780
Araneae, dispersal, 1738
  recolonization, 1738
archytas marmoratus, biological control, 1122
  Heliothis zea, 1122
Artogeia rapae, enzyme-linked immunosorbent assay, 111
  epicuticular wax, 730
  granulosis virus, 121
  Plutella xylostella, 730
Atrichopogon spp., Ceratopogonidae, 286
  Pistia stratiotes, 286
attractants, *Acalymma vittatum*, 8
  Anastrepha ludens, 403
  bacteria, 1726
  Dacus dorsalis, 1726
  insect–plant interactions, 8
  Sargentia greggii, 403
attraction, *Gossypium*, 490
  Pectinophora gossypiella, 490
Aulonium ruficorne, biological control, 1010
  Orthotomicus erosus, 1010

B
Baccharis salicifolia, ecology, 515
  flower feeders, 515
Bacillus thuringiensis, *Chrysomela scripta*, 428
  Ostrinia nubilalis, 182
  phagostimulants, 182
  Populus spp., 428
bacteria, attractants, 1726
  Dacus dorsalis, 1726
baculovirus, granulosis virus, 791
  Plodia interpunctella, 791
baculoviruses, microbial control, 1115
  Oryctes rhinoceros, 1115
Beauveria bassiana, aphid-pathogenic fungi, 815
  bioassay, 815
  biological control, 785
  Diuraphis noxia, 785
bees, crop pollination, 1649
  management systems, 1649
behavior, biology, 44
  Cochliomyia hominivorax, 808
  Empoasca fabae, 524
  reproduction, 808
  Solenopsis invicta, 44
  Sorghum bicolor, 1798
  vegetables, 524
Betula spp., *Lymantria dispar*, 937
  water stress, 937
bioassay, aphid–pathogenic fungi, 815
  Beauveria bassiana, 815
biological control, *Aphidius liriodendrii*, 1130
  Aphidius nigripes, 222
  archytas marmoratus, 1122
  Aulonium ruficorne, 1010
  Beauveria bassiana, 785
  braconid parasites, 1155
  Centaurea solstitialis, 1882
  Ceratitis capitata, 383
  Coleophora laricella, 1873
  Delia antiqua, 1163
  Diatraea grandiosella, 1155, 1889
  Diuraphis noxia, 785
  Elasmopalpus lignosellus, 137
  Endopiza viteana, 764
  epigeal predators, 309
  foraging, 222
  Heliothis zea, 1122
  insect pathogens, 137
  leafminer parasitoids, 625
  Liriomyza spp., 625
  Lymantria dispar, 943
  marigolds, 825
  Melipotis indomita, 415
  Mimosa pigra, 776
  Musca domestica, 578
  Musca domestica, 322
  Nasonia vitripennis, 322
  natural enemy evaluation, 1889
  Neohydronomus affinis, 799
  Ooencyrtus kuvanae, 943
  Orthotomicus erosus, 1010
  parasitic nematodes, 383
  parasitism, 409
  parasitoids, 764, 825, 1873
  phytophagous insects, 776
  Pistia stratiotes, 799
  Prosopis spp., 415
  relative humidity, 1163

Rhagoletis pomonella, 309
Sesbania punicea, 1866
Trichapion lativentre, 1866
Trichogramma spp., 154
Trioxys curvicaudus, 1130
urban entomology, 409
weeds, 1882
biology, behavior, 44
Solenopsis invicta, 44
Sphenophorus spp., 1037
taxonomy, 1037
biometrics, *Diatraea saccharalis*, 635
fecundity, 635
biotypes, leafhoppers, 167
biotypes, *Mayetiola destructor*, 494
biotypes, *Myzus persicae*, 32
Nezara spp., 387
plant resistance, 494
polymorphism, 32
rice, 167
Trissolcus spp., 387
Biprorulus bibax, developmental rates, 874
developmental thresholds, 874
phenology, 1710
reproductive status, 1710
black locust, *Megacyllene robiniae*, 501
tunnel distribution, 501
black spruce, *Choristoneura fumiferana*, 257
egg mass size, 257
body size, *Culex nigripalpus*, 469
rainfall, 469
Boophilus microplus, microclimate, 66
survival, 66
Brachycorynella asparagi, *Diaeretiella rapae*, 890
temperature, 890
braconid parasites, biological control, 1155
Diatraea grandiosella, 1155
Brassicaceae, host plant location, 704
Plutella xylostella, 704
Bupalus piniarius, *Panolis flammea*, 481
Pinus sylvestris, 481

C

cabbage, host plant resistance, 1086
Plutella xylostella, 1086
Cacopsylla pyricola, diapause, 544, 551
mating behavior, 551
oviposition, 357
pears, 357
photoperiod, 544
egg survival, 656
oviposition, 656
Calocoris norvegicus, development model, 848
pistachios, 848
Campoletis sonorensis, mate location, 26
parasitoids, 26
capture rates, *Homadaula anisocentra*, 1
sex peromones, 1
Carabidae, *Listronotus oregonensis*, 1146
voracity and food preference, 1146
cassavas, dry matter, 1767
Mononychellus tanajoa, 1767
Centaurea solstitialis, biological control, 1882
weeds, 1882
Ceratitis capitata, biological control, 383
mortality, 1666
ovipositional preference, 1075
parasitic nematodes, 383
predation, 148
rank order of host choice, 1075
soil effects, 1666
Solenopsis geminata, 148
Ceratopogonidae, *Atrichopogon* spp., 286
Pistia stratiotes, 286
Chironomus decorus, life history, 1681
thermal tolerance, 1681
Choristoneura fumiferana, black spruce, 257
egg mass size, 257
Chrysomela scripta, *Bacillus thuringiensis*, 428
Populus spp., 428
Chrysomelidae, specificity, 905
willows, 905
Chrysoteuchia topiaria, monitoring plant preference, 721
climate, evaluation, 130
weed biocontrol agents, 130
Coccinellidae, hibernation, 1657
lipids, 1657
Cochliomyia hominivorax, behavior, 808
reproduction, 808
Coleophora laricella, biological control, 1873
parasitoids, 1873
communities, Acrididae, 1756
grazing history, 1756
community structure, lindane, 898
species diversity, 898
competition, *Matsucoccus resinosae*, 672
Pineus boerneri, 672
conidial germination, fungal pathogen, 85
Lymantria dispar, 85
conservation tillage, Anthomyiidae, 510
Glycine max, 510
Coptotermes formosanus, *Reticulitermes flavipes*, 15
water relations, 15
corn, adaptation, 949
Diatraea grandiosella, 684
key-factor analysis, 684
Schizaphis graminum, 949
Cotesia melanoscela, *Lymantria dispar*, 771, 859
parasitism, 771
temperature-dependent growth, 859
cotton, *Anthonomus grandis grandis*, 1110
Heliothis spp., 710
oviposition, 710
parasitoids, 1110
courtship behavior, mating competitiveness, 21
Pectinophora gossypiella, 21
cowpeas, *Maruca testulis*, 1792
resistance, 1792
crab apples, plant resistance, 565
Rhagoletis pomonella, 565
crested wheatgrass, predators, 244
sap feeders, 244
crop pollination, bees, 1649
management systems, 1649
Cryphonectria (Endothia) parasitica, *Agrilus bilineatus*, 239
tree stress, 239
Culex nigripalpus, body size, 469
rainfall, 469
Culicoides spp., *Dasyhelea* spp., 1068
Leptoconop spp., 1068
cultural control, *Acigona ignefusalis*, 558
population ecology, 558
Cydia pomonella, monitoring, 573
trap location, 573
Cylas formicarius elegantulus, *Ipomoea batatas*, 313
oviposition, 1663
population dynamics, 313
stimulant boehmeryl, 1663

D

Dacnusa dryas, *Agromyza frontella*, 423
encapsulation, 423
Dacus cucurbitae, *Dacus dorsalis*, 1820
population ecology, 1820

Dacus dorsalis, attractants, 1726
bacteria, 1726
Dacus cucurbitae, 1820
population ecology, 1820
daily mortality, *Lymantria dispar*, 448
simulation, 448
Danthonia parryi, *Festuca scabrella*, 1103
grasshoppers, 1103
Dasyhelea spp., *Culicoides* spp., 1068
Leptoconop spp., 1068
degree-day accumulation, *Manduca* spp., 1055
ovipositional trends, 1055
degree-day survival, *Anticarsia gemmatalis*, 1017
models, 1017
Delia antiqua, biological control, 1163
relative humidity, 1163
Dendroctonus frontalis, geographical variation, 1671
pheromones, 1671
density correlation, Acrididae, 1786
weather, 1786
detoxication, esterases, 842
Lymantria dispar, 842
detrended correspondence analysis, nontarget organisms, 1746
Tenebrionidae, 1746
development, *Graminella nigrifrons*, 209
temperature, 209
development model, *Calocoris norvegicus*, 848
pistachioe, 848
development rates, *Harrisina brillians*, 1689
Spissistilus festinus, 229
temperature effects, 229, 1689
developmental rates, *Biprorulus bibax*, 874
developmental thresholds, 874
developmental thresholds, *Biprorulus bibax*, 874
developmental rates, 874
Diabrotica barberi, *Diabrotica virgifera virgifera*, 882
semiochemicals, 882
Diabrotica undecimpunctata howardi, predator-weed interactions, 1837
soil moisture, 697
soil texture, 697
weeds, 1837
Diabrotica virgifera virgifera, *Diabrotica barberi*, 882
larval movement, 666
semiochemicals, 882
soil, 666
Diabrotica spp., insect-plant relationships, 474
Diaeretiella rapae, *Brachycorynella asparagi*, 890
temperature, 890
diapause, *Ancylis comptana*, 932
Anthonomus grandis grandis, 91
Cacopsylla pyricola, 544
Cacopsylla pyricola, 551
Diatraea spp., 345
dormancy, 932
Eoreuma loftini, 345
mating behavior, 551
mortality, 91
photoperiod, 544
Diatraea grandiosella, biological control, 1889
biological control, 1155
braconid parasites, 1155
corn, 684
key-factor analysis, 684
natural enemy evaluation, 1889
Diatraea saccharalis, biometrics, 635
fecundity, 635
Diatraea spp., diapause, 345
Eoreuma loftini, 345
Eoreuma spp., 393
parasites, 393
dispersal, *Anopheles quadrimaculatus*, 961
Anthonomus grandis grandis, 195
Araneae, 1738
flight initiation, 281
flight range, 961
Leptinotarsa decemlineata, 281
migration, 1829
modeling, 195
pine root weevils, 1829
Platynota idaeusalis, 648
recolonization, 1738
rubidium, 648
dispersal agents, *Adelges tsugae*, 36
insect dispersal, 36
Diuraphis noxia, *Beauveria bassiana*, 785
biological control, 785
dormancy, *Ancylis comptana*, 932
diapause, 932
dry matter, cassavas, 1767
Mononychellus tanajoa, 1767

E

ecology, *Baccharis salicifolia*, 515
flower feeders, 515
ectoparasite, Lepidoptera, 795
nematoda, 795
Edovum puttleri, honeydews, 1732
Leptinotarsa decemlineata, 1732
egg development, *Neodiprion fulviceps*, 456
oviposition, 456
egg mass length, fecundity, 108
Lymantria dispar, 108
egg mass size, black spruce, 257
Choristoneura fumiferana, 257
egg parasitism, *Heliothis* spp., 753
Trichogramma spp., 753
egg survival, *Cacopsylla pyricola*, 656
oviposition, 656
Elasmopalpus lignosellus, biological control, 137
insect pathogens, 137
Empoasca fabae, behavior, 524
intercropping, 534
Empoasca fabae, *Medicago sativa*, 176
photosynthesis, 176
population dynamics, 534
relay intercropping, 1810
soybeans, 1810
vegetables, 524
Empoasca spp., parasitoids, 505
predation, 505
encapsulation, *Agromyza frontella*, 423
Dacnusa dryas, 423
endocrine regulation, *Leptinotarsa decemlineata*, 117
migration, 117
Endopiza viteana, biological control, 764
parasitoids, 764
environment, flight activity, 293
Winthemia fumiferanae, 293
environmental chambers, *Pseudaletia unipuncta*, 866
thermoperiod, 866
enzyme-linked immunosorbent assay, *Artogeia rapae*, 111
granulosis virus, 111
Eoreuma loftini, diapause, 345
Diatraea spp., 345
Eoreuma spp., *Diatraea* spp., 393
parasites, 393
epicuticular wax, *Artogeia rapae*, 730
Plutella xylostella, 730
epigeal predators, biological control, 309
Rhagoletis pomonella, 309
Epilachna varivestis, induced resistance, 1852
induced resistance, 854
soybeans, 854, 1852
Epiphyas postvittana, mating disruption, 1702
pheromones, 1702

Erythroneura spp., *Vitis vinifera*, 1803
water stress, 1803
esterases, detoxication, 842
Lymantria dispar, 842
Etiella zinckenella, host plant density, 640
soybeans, 640
Eurosta, parasitoids, 972
Solidago spp., 972
evaluation, climate, 130
weed biocontrol agents, 130

F
fecundity, biometrics, 635
Diatraea saccharalis, 635
egg mass length, 108
Lymantria dispar, 108, 1024
rhythm, 630
sampling, 1024
Thyanta pallidovirens, 630
feeding behavior, *Nephotettix virescens*, 594
resistance, 594
Festuca scabrella, *Danthonia parryi*, 1103
grasshoppers, 1103
fire ants, *Solenopsis invicta*, 878
temperature, 878
flight activity, environment, 293
Winthemia fumiferanae, 293
flight initiation, dispersal, 281
Leptinotarsa decemlineata, 281
flight range, *Anopheles quadrimaculatus*, 961
dispersal, 961
flower feeders, *Baccharis salicifolia*, 515
ecology, 515
flush bar technique, action site, 1081
Ostrinia nubilalis, 1081
foraging, *Aphidius nigripes*, 222
biological control, 222
fruit, symbionts, 995
yeast, 995
fungal pathogens, conidial germination, 85
Hypera postica, 612
Lymantria dispar, 85
phenology, 612

G
geographic information systems, *Lymantria dispar*, 370
population dynamics, 370
geographical variation, *Dendroctonus frontalis*, 1671
pheromones, 1671
Glycine max, Anthomyiidae, 510
conservation tillage, 510
Gossypium, attraction, 490
Pectinophora gossypiella, 490
Gossypium hirsutum, *larval distribution*, *1060*
Spodoptera frugiperda, 1060
Gossypium spp., *Apis mellifera*, 1034
pollen deposition, 1034
gossyplure, *Pectinophora gossypiella*, 1845
pheromones, 1845
Graminella nigrifrons, development, 209
temperature, 209
granulosis virus, *Artogeia rapae*, 111
baculovirus, 791
enzyme-linked immunosorbent assay, 111
Plodia interpunctella, 791
grasses, leafhoppers, 76
maize chlorotic dwarf virus, 76
grasshoppers, *Danthonia parryi*, 1103
Festuca scabrella, 1103
grazing history, Acrididae, 1756
communities, 1756

H
Harrisina brillians, development rates, 1689
temperature effects, 1689
Helicoverpa armigera, parasitism, 1141
parasitoid-crop associations, 1141
Heliothis spp., cotton, 710
egg parasitism, 753
oviposition, 710
Trichogramma spp., 753
Heliothis virescens, *Microplitis croceipes*, 251
nuclear polyhedrosis virus, 251
Heliothis zea, *archytas marmoratus*, 1122
biological control, 1122
parasitoids, 1135
population dynamics, 274
soybeans, 1135
Hemadas nubilipennis, lowbush blueberry galls, 911
parasitoid community, 911
herbs, antifeedants, 234
Leptinotarsa decemlineata, 234
hibernation, Coccinellidae, 1657
lipids, 1657
Homadaula anisocentra, capture rates, 1
sex peromones, 1
honeydews, *Edovum puttleri*, 1732
Leptinotarsa decemlineata, 1732
host finding, pheromones, 1675
Smicronyx fulus, 1675
host habitat, *Papaipema nebris*, 162
parasitism, 162
host plant density, *Etiella zinckenella*, 640
soybeans, 640
host plant location, Brassicaceae, 704
Plutella xylostella, 704
host plant resistance, cabbage, 1086
Ostrinia nubilalis, 1091
Plutella xylostella, 1086
Zea mays, 1091
host preference parasitoid, *Pholetesor ornigis*, 1097
Phyllonorycter blancardella, 1097
host specificity, karyotypes, 609
Rhopalosiphum maidis, 609
host stage preferences, parasitoid-induced mortality, 803
Pnigalio flavipes, 803
host-finding behavior, oviposition behavior, 142
parasites, 142
Hypera postica, fungal pathogens, 612
phenology, 612

I
induced resistance, *Epilachna varivestis*, 854
Epilachna varivestis, 1852
soybeans, 854, 1852
insect dispersal, *Adelges tsugae*, 36
dispersal agents, 36
Popillia japonica, 955
strip-cropping, 955
insect pathogens, biological control, 137
Elasmopalpus lignosellus, 137
insect-plant interactions, *Acalymma vittatum*, 8
attractants, 8
Plutella xylostella, 587
reproductive maturation, 587
insect-plant relationships, *Diabrotica* spp., 474
Mayetiola destructor, 303
oviposition, 303
insecticide toxicity, pest management, 439
population models, 439
insecticides, *Musca domestica*, 1722
synergists, 1722
integrated pest management, *Ancylis comptana*, 339
temperature-development relationship, 339
intercropping, *Empoasca fabae*, 534
population dynamics, 534
interplanting, oviposition, 987
Plutella xylostella, 987

iodoacetic acid, soybeans, 215
Trichoplusia ni, 215
Ipomoea batatas, *Cylas formicarius elegantulus*, 313
population dynamics, 313
Ixodes dammini, larval dispersal, 1029
Lyme disease, 1029

K
karyotypes, host specificity, 609
Rhopalosiphum maidis, 609
key-factor analysis, corn, 684
Diatraea grandiosella, 684

L
larval dispersal, *Ixodes dammini*, 1029
Lyme disease, 1029
larval distribution, *Gossypium hirsutum*, 1060
Spodoptera frugiperda, 1060
larval movement, *Diabrotica virgifera virgifera*, 666
soil, 666
latitudinal variation, *Acyrthosiphon pisum*, 618
photoperiod, 618
leafhoppers, biotypes, 167
grasses, 76
maize chlorotic dwarf virus, 76
rice, 167
leafminer parasitoids, biological control, 625
Liriomyza spp., 625
Lepidoptera, ectoparasite, 795
nematoda, 795
Leptinotarsa decemlineata, antifeedants, 234
dispersal, 281
Edovum puttleri, 1732
endocrine regulation, 117
flight initiation, 281
herbs, 234
honeydews, 1732
life history strategy, 123
migration, 117
parasites, oviposition, 104
r-strategists, 123
Leptoconop spp., *Culicoides* spp., 1068
Dasyhelea spp., 1068
life history, *Chironomus decorus*, 1681
thermal tolerance, 1681
life history strategy, *Leptinotarsa decemlineata*, 123
r-strategists, 123
life table, *Acyrthosiphon kondoi*, 1001
natural control, 1001
lindane, community structure, 898
species diversity, 898
lipids, Coccinellidae, 1657
hibernation, 1657
Liriomyza spp., biological control, 625
leafminer parasitoids, 625
Listronotus oregonensis, *Carabidae*, 1146
reproductive diapause, 836
temperature–photoperiod regimens, 836
voracity and food preference, 1146
Lonchaea corticis, *Picea sitchensis*, 54
Pissodes strobi, 54
lowbush blueberry galls, *Hemadas nubilipennis*, 911
parasitoid community, 911
Lymantria dispar, *Betula* spp., 937
biological control, 943
conidial germination, 85
Cotesia melanoscela, 771, 859
daily mortality, 448
detoxication, 842
egg mass length, 108
esterases, 842
fecundity, 108, 1024
fungal pathogen, 85
geographic information systems, 370
nuclear polyhedrosis virus, 662
Ooencyrtus kuvanae, 943
parasitism, 771
population dynamics, 370
sampling, 1024
simulation, 448
temperature-dependent growth, 859
transmission, 662
water stress, 937
Lyme disease, *Ixodes dammini*, 1029
larval dispersal, 1029

M
maize chlorotic dwarf virus, grasses, 76
leafhoppers, 76
management systems, bees, 1649
crop pollination, 1649
Manduca spp., degree-day accumulation, 1055
ovipositional trends, 1055
marigolds, biological control, 825
parasitoids, 825
Maruca testulis, cowpeas, 1792
resistance, 1792
mate location, *Campoletis sonorensis*, 26
parasitoids, 26
mating behavior, *Cacopsylla pyricola*, 551
diapause, 551
mating competitiveness, courtship behavior, 21
Pectinophora gossypiella, 21
mating disruption, *Epiphyas postvittana*, 1702
pheromones, 1702
Matsucoccus resinosae, competition, 672
Pinus boerneri, 672
Mayetiola destructor, biotypes, 494
insect-plant relationships, 303
oviposition, 303
plant resistance, 494
soluble carbohydrates, 1698
wheat plant resistance, 1698
Medicago sativa, *Empoasca fabae*, 176
photosynthesis, 176
Megacyllene robiniae, black locust, 501
tunnel distribution, 501
Melipotis indomita, biological control, 415
Prosopis spp., 415
microbial control, baculoviruses, 1115
Oryctes rhinoceros, 1115
microclimate, *Boophilus microplus*, 66
Phlebotomus spp., 486
Sergentomyia spp., 486
survival, 66
microhabitat distribution, *Anticarsia gemmatalis*, 740
soybean, 740
Microplitis croceipes, *Heliothis virescens*, 251
nuclear polyhedrosis virus, 251
migration, dispersal, 1829
endocrine regulation, 117
Leptinotarsa decemlineata, 117
Peridroma saucia, 603
phenology, 603
pine root weevils, 1829
Mimosa pigra, biological control, 776
phytophagous insects, 776
modeling, *Anthonomus grandis grandis*, 195
dispersal, 195
models, *Anticarsia gemmatalis*, 1017
degree-day survival, 1017
phenology, 916
Quadraspidiotus perniciosus, 916
monitoring, *Cydia pomonella*, 573
trap location, 573
monitoring plant preference,

Chrysoteuchia topiaria, 721
Mononychellus tanajoa, cassavas, 1767
dry matter, 1767
mortality, *Anthonomus grandis grandis*, 59, 91
Ceratitis capitata, 1666
diapause, 91
nuclear polyhedrosis virus, 1150
parasitism, 59
soil effects, 1666
Spodoptera ornithogalli, 1150
multiple regression, nuclear polyhedrosis, 260
Spodoptera frugiperda, 260
Musca domestica, biological control, 322, 578
insecticides, 1722
Nasonia vitripennis, 322
synergists, 1722
Myzus persicae, biotypes, 32
polymorphism, 32

N
Nasonia vitripennis, biological control, 322
Musca domestica, 322
natural control, *Acyrthosiphon kondoi*, 1001
life table, 1001
natural enemy evaluation, biological control, 1889
Diatraea grandiosella, 1889
nematoda, ectoparasite, 795
Lepidoptera, 795
Neodiprion fulviceps, egg development, 456
oviposition, 456
Neohydronomus affinis, biological control, 799
Pistia stratiotes, 799
Nephotettix virescens, feeding behavior, 594
resistance, 594
Nezara spp., biotype, 387
Trissolcus spp., 387
nontarget organisms, detrended correspondence analysis, 1746
Tenebrionidae, 1746
nuclear polyhedrosis, multiple regression, 260
Spodoptera frugiperda, 260
nuclear polyhedrosis virus, *Heliothis virescens*, 251
Lymantria dispar, 662
Microplitis croceipes, 251
mortality, 1150
Spodoptera ornithogalli, 1150
transmission, 662

O
onion nectar, *Allium cepa*, 327
Apis mellifera, 327
Oencyrtus kuvanae, biological control, 943
Lymantria dispar, 943
Oracella acuta, pine seed orchards, 99
population densities, 99
Orthotomicus erosus, *Aulonium ruficorne*, 1010
biological control, 1010
Oryctes rhinoceros, baculoviruses, 1115
microbial control, 1115
Osmia lignaria propinqua, pollen, 332
polylecty, 332
Ostrinia nubilalis, action site, 1081
Bacillus thuringiensis, 182
flush bar technique, 1081
host plant resistance, 1091
phagostimulants, 182
sweet corn, 1858
weed–insect interactions, 1858
Zea mays, 1091
oviposition, *Cacopsylla pyricola*, 357, 656
cotton, 710
Cylas formicarius elegantulus, 1663
egg development, 456
egg survival, 656
Heliothis spp., 710
insect–plant relationships, 303
interplanting, 987
Leptinotarsa decemlineata parasites, 104
Mayetiola destructor, 303
Neodiprion fulviceps, 456
pear, 357
Plutella xylostella, 987
stimulant boehmeryl, 1663
oviposition behavior, host-finding behavior, 142
parasites, 142
ovipositional preference, *Ceratitis capitata*, 1075
rank order of host choice, 1075
soil moisture, 966
southern corn rootworms, 966
ovipositional trends, degree-day accumulation, 1055
Manduca spp., 1055

P
Panolis flammea, *Bupalus piniarius*, 481
Pinus sylvestris, 481
Papaipema nebris, host habitat, 162
parasitism, 162
parasites, *Diatraea* spp., 393
Eoreuma spp., 393
host-finding behavior, 142
oviposition behavior, 142
parasitic nematodes, biological control, 383
Ceratitis capitata, 383
parasitism, *Anthonomus grandis grandis*, 59
biological control, 409
Cotesia melanoscela, 771
Helicoverpa armigera, 1141
host habitat, 162
Lymantria dispar, 771
mortality, 59
Papaipema nebris, 162
parasitoid–crop associations, 1141
phenology, 362
Phyllonorycter elmaella, 362
urban entomology, 409
parasitoid community, *Hemadas nubilipennis*, 911
lowbush blueberry galls, 911
parasitoid–crop associations, *Helicoverpa armigera*, 1141
parasitism, 1141
parasitoid–induced mortality, host stage preferences, 803
Pnigalio flavipes, 803
parasitoids, *Anthonomus grandis grandis*, 1110
Anticarsia gemmatalis, 746
biological control, 764, 825, 1873
Campoletis sonorensis, 26
Coleophora laricella, 1873
cotton, 1110
Empoasca spp., 505
Endopiza viteana, 764
Eurosta, 972
Heliothis zea, 1135
marigolds, 825
mate location, 26
pheromones, 926
predation, 505
Pseudoplusia includens, 746
Quadraspidiotus perniciosus, 926
Solidago spp., 972
soybeans, 1135
pear, *Cacopsylla pyricola*, 357
oviposition, 357
Pectinophora gossypiella, attraction, 490
courtship behavior, 21
Gossypium, 490
gossyplure, 1845
mating competitiveness, 21
pheromones, 1845
Peridroma saucia, migration, 603
phenology, 603
pest management, insecticide toxicity, 439
population models, 439

phagostimulants, *Bacillus thuringiensis*, 182
Ostrinia nubilalis, 182
phenology, *Biprorulus bibax*, 1710
fungal pathogens, 612
Hypera postica, 612
migration, 603
models, 916
parasitism, 362
Peridroma saucia, 603
Phyllonorycter elmaella, 362
Quadraspidiotus perniciosus, 916
reproductive status, 1710
pheromones, *Dendroctonus frontalis*, 1671
Epiphyas postvittana, 1702
geographical variation, 1671
gossyplure, 1845
host finding, 1675
mating disruption, 1702
parasitoids, 926
Pectinophora gossypiella, 1845
Quadraspidiotus perniciosus, 926
Smicronyx fulus, 1675
Phlebotomus spp., microclimate, 486
Sergentomyia spp., 486
Pholetesor ornigis, host preference parasitoid, 1097
Phyllonorycter blancardella, 1097
photoperiod, *Acyrthosiphon pisum*, 618
Cacopsylla pyricola, 544
diapause, 544
latitudinal variation, 618
photosynthesis, *Empoasca fabae*, 176
Medicago sativa, 176
Phyllocoptruta oleivora, *Polyphagotarsonemus latus*, 378
spatial distribution, 378
Phyllonorycter blancardella, host preference parasitoid, 1097
Pholetesor ornigis, 1097
Phyllonorycter elmaella, parasitism, 362
phenology, 362
phytophagous insects, biological control, 776
Mimosa pigra, 776
Picea sitchensis, *Lonchaea corticis*, 54
Pissodes strobi, 54
pine root weevils, dispersal, 1829
migration, 1829
pine seed orchards, *Oracella acuta*, 99
population densities, 99
Pineus boerneri, competition, 672
Matsucoccus resinosae, 672
Pinus sylvestris, *Bupalus piniarius*, 481
Panolis flammea, 481
Pissodes strobi, *Lonchaea corticis*, 54
Picea sitchensis, 54
pistachios, *Calocoris norvegicus*, 848
development model, 848
Pistia stratiotes, *Atrichopogon* spp., 286
biological control, 799
Ceratopogonidae, 286
Neohydronomus affinis, 799
plant chemistry, *Solidago altissima*, 465
Trichoplusia ni, 465
plant resistance, biotypes, 494
crab apples, 565
Mayetiola destructor, 494
Rhagoletis pomonella, 565
Platynota idaeusalis, dispersal, 648
rubidium, 648
Plodia interpunctella, baculovirus, 791
granulosis virus, 791
Plutella xylostella, *Artogeia rapae*, 730
Brassicaceae, 704
cabbage, 1086
epicuticular wax, 730
host plant location, 704
host plant resistance, 1086
insect–plant interaction, 587
interplanting, 987
oviposition, 987
reproductive maturation, 587
Pnigalio flavipes, host stage preferences, 803
parasitoid–induced mortality, 803
pollen, *Osmia lignaria propinqua*, 332
polylecty, 332
pollen deposition, *Apis mellifera*, 1034
Gossypium spp., 1034
polylecty, *Osmia lignaria propinqua*, 332
pollen, 332
polymorphism, biotypes, 32
Myzus persicae, 32
Polyphagotarsonemus latus, *Phyllocoptruta oleivora*, 378
spatial distribution, 378
Popillia japonica, insect dispersal, 955
strip–cropping, 955
population densities,-*Oracella acuta*, 99
pine seed orchards, 99
population dynamics, *Cylas formicarius elegantulus*, 313
Empoasca fabae, 534
geographic information systems, 370
Heliothis zea, 274
intercropping, 534
Ipomoea batatas, 313
Lymantria dispar, 370
soybean pests, 1773
Tetranychus urticae, 1773
population ecology, *Acigona ignefusalis*, 558
cultural control, 558
Dacus cucurbitae, 1820
Dacus dorsalis, 1820
population models, insecticide toxicity, 439
pest management, 439
Populus spp., *Bacillus thuringiensis*, 428
Chrysomela scripta, 428
predation, *Ceratitis capitata*, 148
Empoasca spp., 505
parasitoids, 505
Solenopsis geminata, 148
predator–weed interactions, *Diabrotica undecimpunctata howardi*, 1837
weeds, 1837
predators, crested wheatgrass, 244
sap feeders, 244
Prosopis spp., biological control, 415
Melipotis indomita, 415
Pseudaletia unipuncta, environmental chambers, 866
thermoperiod, 866
Pseudoplusia includens, *Anticarsia gemmatalis*, 717, 746
sprinkler irrigation, 717

Q

Quadraspidiotus perniciosus, models, 916
parasitoids, 926
phenology, 916
pheromones, 926

R

r–strategists, *Leptinotarsa decemlineata*, 123
life history strategy, 123
rainfall, body size, 469
Culex nigripalpus, 469
rank order of host choice, *Ceratitis capitata*, 1075
ovipositional preference, 1075
recolonization, Araneae, 1738
dispersal, 1738
relative humidity, biological control, 1163
Delia antiqua, 1163

relay intercropping, *Empoasca fabae*, 1810
soybeans, 1810
reproduction, behavior, 808
Cochliomyia hominivorax, 808
reproductive diapause, *Listronotus oregonensis*, 836
temperature–photoperiod regimens, 836
reproductive maturation, insect–plant interaction, 587
Plutella xylostella, 587
reproductive status, *Biprorulus bibax*, 1710
phenology, 1710
resistance, cowpeas, 1792
feeding behavior, 594
Maruca testulis, 1792
Nephotettix virescens, 594
Rhyzopertha dominica, 268
Sitotroga cerealella, 268
Reticulitermes flavipes, *Coptotermes formosanus*, 15
water relations, 15
Rhagoletis pomonella, crab apples, 565
plant resistance, 565
Rhagoletis pomonella, biological control, 309
epigeal predators, 309
Rhopalosiphum maidis, host specificity, 609
karyotypes, 609
rhythm, fecundity, 630
Thyanta pallidovirens, 630
Rhyzopertha dominica, resistance, 268
simulation models, 725
Sitotroga cerealella, 268
stored products, 725
rice, biotypes, 167
leafhoppers, 167
robber flies, *Stenopogon inquinatus*, 459
thermoregulation, 459
rubidium, dispersal, 648
Platynota idaeusalis, 648

S

sampling, fecundity, 1024
Lymantria dispar, 1024
sap feeders, crested wheatgrass, 244
predators, 244
Sargentia greggii, *Anastrepha ludens*, 403
attractants, 403
Schizaphis graminum, adaptation, 949
corn, 949
semiochemicals, *Diabrotica barberi*, 882
Diabrotica virgifera virgifera, 882
Sergentomyia spp., microclimate, 486
Phlebotomus spp., 486
Sesbania punicea, biological control, 1866
Trichapion lativentre, 1866
sex peromones, capture rates, 1
Homadaula anisocentra, 1
sexual receptivity, *Supella longipalpa* mating, 1716
simulation, daily mortality, 448
Lymantria dispar, 448
simulation models, *Rhyzopertha dominica*, 725
stored products, 725
Sitotroga cerealella, resistance, 268
Rhyzopertha dominica, 268
Smicronyx fulus, host finding, 1675
pheromones, 1675
soil, *Diabrotica virgifera virgifera*, 666
larval movement, 666
soil effects, *Ceratitis capitata*, 1666
mortality, 1666
soil moisture, *Diabrotica undecimpunctata howardi*, 697
oviposition preference, 966
soil texture, 697
southern corn rootworms, 966
soil texture, *Diabrotica undecimpunctata howardi*, 697
soil moisture, 697
Solenopsis geminata, *Ceratitis capitata*, 148
predation, 148
Solenopsis invicta, *Anticarsia gemmatalis*, 978
behavior, 44
biology, 44
fire ants, 878
soybeans, 978
temperature, 878
Solidago altissima, plant chemistry, 465
Trichoplusia ni, 465
Solidago spp., *Eurosta*, 972
parasitoids, 972
soluble carbohydrates, *Mayetiola destructor*, 1698
wheat plant resistance, 1698
Sorghum bicolor, behavior, 1798
southern corn rootworms, oviposition preference, 966
soil moisture, 966
soybean pests, population dynamics, 1773
Tetranychus urticae, 1773
soybeans, alfalfa, 677
Anticarsia gemmatalis, 740, 978
Empoasca fabae, 1810
Epilachna varivestis, 854, 1852
Etiella zinckenella, 640
Heliothis zea, 1135
host plant density, 640
induced resistance, 854, 1852
iodoacetic acid, 215
microhabitat distribution, 740
parasitoids, 1135
relay intercropping, 1810
Solenopsis invicta, 978
Spissistilus festinus, 677
Trichoplusia ni, 215
spatial distribution, *Phyllocoptruta oleivora*, 378
Polyphagotarsonemus latus, 378
species diversity, community structure, 898
lindane, 898
specificity, Chrysomelidae, 905
willows, 905
Sphenophorus spp., biology, 1037
taxonomy, 1037
Spissistilus festinus, alfalfa, 677
development rate, 229
soybean, 677
temperature effects, 229
Spodoptera frugiperda, *Gossypium hirsutum*, 1060
larval distribution, 1060
multiple regression, 260
nuclear polyhedrosis, 260
Spodoptera ornithogalli, mortality, 1150
nuclear polyhedrosis virus, 1150
sprinkler irrigation, *Anticarsia gemmatalis*, 717
Pseudoplusia includens, 717
Stenopogon inquinatus, robber flies, 459
thermoregulation, 459
stimulant boehmeryl, *Cylas formicarius elegantulus*, 1663
oviposition, 1663
stored products, *Rhyzopertha dominica*, 725
simulation models, 725
strip–cropping, insect dispersal, 955
Popillia japonica, 955
subterranean termites, agonistic behavior, 1047
territory size, 1047
Supella longipalpa mating, sexual receptivity, 1716
survival, *Boophilus microplus*, 66
microclimate, 66
sweet corn, *Ostrinia nubilalis*, 1858
weed–insect interactions, 1858
symbionts, fruit, 995
yeast, 995

synergists, insecticides, 1722
Musca domestica, 1722

T
taxonomy, biology, 1037
Sphenophorus spp., 1037
temperature, *Apis mellifera*, 1780
Brachycorynella asparagi, 890
development, 209
Diaeretiella rapae, 890
fire ants, 878
Graminella nigrifrons, 209
Solenopsis invicta, 878
Varroa jacobsoni, 1780
temperature effects, development rate, 229, 1689
Harrisina brillians, 1689
Spissistilus festinus, 229
temperature-dependent growth, *Cotesia melanoscela*, 859
Lymantria dispar, 859
temperature-development relationship, *Ancylis comptana*, 339
integrated pest management, 339
temperature-photoperiod regimens, *Listronotus oregonensis*, 836
reproductive diapause, 836
Tenebrionidae, detrended correspondence analysis, 1746
nontarget organisms, 1746
territory size, agonistic behavior, 1047
subterranean termites, 1047
Tetranychus urticae, population dynamics, 1773
soybean pests, 1773
thermal tolerance, *Chironomus decorus*, 1681
life history, 1681
thermoperiod, environmental chambers, 866
Pseudaletia unipuncta, 866
thermoregulation, robber flies, 459
Stenopogon inquinatus, 459
Thyanta pallidovirens, fecundity, 630
rhythm, 630
transmission, *Lymantria dispar*, 662
nuclear polyhedrosis virus, 662
trap location, *Cydia pomonella*, 573
monitoring, 573
tree stress, *Agrilus bilineatus*, 239
Cryphonectria (Endothia) parasitica, 239
Trichapion lativentre, biological control, 1866
Sesbania punicea, 1866
Trichogramma spp., biological control, 154
egg parasitism, 753
Heliothis spp., 753
Trichoplusia ni, iodoacetic acid, 215
plant chemistry, 465
Solidago altissima, 465
soybeans, 215
Trioxys curvicaudus, *Aphidius liriodendrii*, 1130
biological control, 1130
Trissolcus spp., biotype, 387
Nezara spp., 387
tunnel distribution, black locust, 501
Megacyllene robiniae, 501

U
urban entomology, biological control, 409
parasitism, 409

V
Varroa jacobsoni, *Apis mellifera*, 1780
temperatures, 1780
vegetables, behavior, 524
Empoasca fabae, 524
Vitis vinifera, *Erythroneura* spp., 1803
water stress, 1803
voracity and food preference, *Carabidae*, 1146
Listronotus oregonensis, 1146

W
water relations, *Coptotermes formosanus*, 15
Reticulitermes flavipes, 15
water stress, *Betula* spp., 937
Erythroneura spp., 1803
Lymantria dispar, 937
Vitis vinifera, 1803
weather, Acrididae, 1786
density correlation, 1786
weed biocontrol agents, climate, 130
evaluation, 130
weed-insect interactions, *Ostrinia nubilalis*, 1858
sweet corn, 1858
weeds, biological control, 1882
Centaurea solstitialis, 1882
Diabrotica undecimpunctata howardi, 1837
predator-weed interactions, 1837
wheat plant resistance, *Mayetiola destructor*, 1698
soluble carbohydrates, 1698
willows, Chrysomelidae, 905
specificity, 905
Winthemia fumiferanae, environment, 293
flight activity, 293

Y
yeast, fruit, 995
symbionts, 995

Z
Zea mays, host plant resistance, 1091
Ostrinia nubilalis, 1091